

TEMA 12 – CÁLCULO DE PRIMITIVAS

EJERCICIO : Calcula las integrales:

- | | | | |
|---|-------------------------------------|-------------------------------------|-------------------------------------|
| a) $\int \frac{4x - 2 \cos x}{x^2 - \sin x} dx$ | b) $\int x e^{2x} dx$ | c) $\int \frac{x}{x^2 - 9} dx$ | d) $\int \frac{e^x}{1 + e^{2x}} dx$ |
| e) $\int \frac{e^x}{1 + (e^x)^2} dx$ | f) $\int \frac{dx}{x^3 + 2x^2 + x}$ | g) $\int \frac{dx}{\sqrt[4]{3x+1}}$ | h) $\int (2x+1) \cos x dx$ |
| i) $\int \frac{2}{x^2 - 1} dx$ | j) $\int 2x(x-1)^2 dx$ | k) $\int (x+2) \sin x dx$ | l) $\int \frac{dx}{x^3 - 3x^2}$ |
| m) $\int \frac{3x + \sqrt{x}}{x^2} dx$ | n) $\int (x+3) e^{x+1} dx$ | ñ) $\int \frac{x}{x^2 + 2x - 8} dx$ | |

Solución:

a) $\int \frac{4x - 2 \cos x}{x^2 - \sin x} dx = \int \frac{2(2x - \cos x)}{x^2 - \sin x} dx = 2 \ln |x^2 - \sin x| + k$

b) $\int x e^{2x} dx$. Integraremos por partes:
$$\begin{cases} u = x \rightarrow du = dx \\ dv = e^{2x} dx \rightarrow v = \frac{1}{2} e^{2x} \end{cases}$$

$$\int x e^{2x} dx = \frac{x}{2} e^{2x} - \int \frac{1}{2} e^{2x} dx = \frac{x}{2} e^{2x} - \frac{1}{4} e^{2x} + k = \left(\frac{x}{2} - \frac{1}{4}\right) e^{2x} + k$$

c) $\int \frac{x}{x^2 - 9} dx$. Descomponemos en fracciones simples:

$$\frac{x}{x^2 - 9} = \frac{x}{(x-3)(x+3)} = \frac{A}{x-3} + \frac{B}{x+3} = \frac{A(x+3) + B(x-3)}{(x-3)(x+3)}$$

- Para $x=3 \rightarrow 3=6A \rightarrow A=\frac{1}{2}$

- Para $x=-3 \rightarrow -3=-6B \rightarrow B=-\frac{1}{2}$

Por tanto: $\int \frac{x}{x^2 - 9} dx = \int \left(\frac{\frac{1}{2}}{x-3} + \frac{-\frac{1}{2}}{x+3} \right) dx = \frac{1}{2} \ln|x-3| + \frac{1}{2} \ln|x+3| + k$

d) $\int \frac{e^x}{1 + e^{2x}} dx = \int \frac{e^x}{1 + (e^x)^2} dx = \arctg(e^x) + k$

e) $\int (x+2) \ln x dx$. Integraremos por partes:
$$\begin{cases} u = \ln x \rightarrow du = \frac{1}{x} dx \\ dv = (x+2) dx \rightarrow v = \frac{x^2}{2} + 2x \end{cases}$$

$$\int (x+2) \ln x dx = \left(\frac{x^2}{2} + 2x \right) \ln x - \int \left(\frac{x^2}{2} + 2x \right) \frac{1}{x} dx = \left(\frac{x^2}{2} + 2x \right) \ln x - \int \left(\frac{x}{2} + 2 \right) dx =$$

$$= \left(\frac{x^2}{2} + 2x \right) \ln x - \frac{x^2}{4} - 2x + k$$

f) $\int \frac{dx}{x^3 + 2x^2 + x}$. Descomponemos en fracciones simples:

$$\frac{1}{x^3 + 2x^2 + x} = \frac{1}{x(x+1)^2} = \frac{A}{x} + \frac{B}{x+1} + \frac{C}{(x+1)^2} = \frac{A(x+1)^2 + Bx(x+1) + Cx}{x(x+1)^2}$$

- Para $x=0 \rightarrow 1=A$
- Para $x=-1 \rightarrow 1=-C \rightarrow C=-1$
- Para $x=1 \rightarrow 1=4A+2B+C \rightarrow B=-1$

Por tanto: $\int \frac{dx}{x^3 + 2x^2 + x} = \int \left(\frac{1}{x} + \frac{-1}{x+1} + \frac{-1}{(x+1)^2} \right) dx = \ln|x| - \ln|x+1| + \frac{1}{x+1} + k$

g) $\int \frac{dx}{\sqrt[4]{3x+1}} = \int (3x+1)^{-1/4} dx = \frac{1}{3} \int 3(3x+1)^{-1/4} dx = \frac{1}{3} \frac{(3x+1)^{3/4}}{\frac{3}{4}} + k = \frac{4\sqrt[4]{(3x+1)^3}}{9} + k$

h) $\int (2x+1)\cos x dx$. Integraremos por partes: $\begin{cases} u = 2x+1 & \rightarrow du = 2dx \\ dv = \cos x dx & \rightarrow v = \sin x \end{cases}$

$$\int (2x+1)\cos x dx = (2x+1)\sin x - \int 2\sin x dx = (2x+1)\sin x + 2\cos x + k$$

i) $\int \frac{2}{x^2 - 1} dx$. Descomponemos en fracciones simples:

$$\frac{2}{x^2 - 1} = \frac{2}{(x-1)(x+1)} = \frac{A}{x-1} + \frac{B}{x+1} = \frac{A(x+1) + B(x-1)}{(x-1)(x+1)}$$

- Para $x=1 \rightarrow 2=2A \rightarrow A=1$
- Para $x=-1 \rightarrow 2=-2B \rightarrow B=-1$

Por tanto: $\int \frac{2}{x^2 - 1} dx = \int \left(\frac{1}{x-1} + \frac{-1}{x+1} \right) dx = \ln|x-1| - \ln|x+1| + k$

j) $\int 2x(x-1)^2 dx = \int 2x(x^2 - 2x + 1) dx = \int (2x^3 - 4x^2 + 2x) dx = \frac{2x^4}{4} - \frac{4x^3}{3} + x^2 + k = \frac{x^4}{2} - \frac{4x^3}{3} + x^2 + k$

k) $\int (x+2)\sin x dx$. Integraremos por partes: $\begin{cases} u = x+2 & \rightarrow du = dx \\ dv = \sin x dx & \rightarrow v = -\cos x \end{cases}$

$$\int (x+2)\sin x dx = -(x+2)\cos x + \int \cos x dx = -(x+2)\cos x + \sin x + k$$

l) $\int \frac{dx}{x^3 - 3x^2}$. Descomponemos en fracciones simples:

$$\frac{1}{x^3 - 3x^2} = \frac{1}{x^2(x-3)} = \frac{A}{x} + \frac{B}{x^2} + \frac{C}{x-3} = \frac{Ax(x-3) + B(x-3) + Cx^2}{x^2(x-3)}$$

- Para $x=0 \rightarrow 1=-3B \rightarrow B=\frac{-1}{3}$

- Para $x=3 \rightarrow 1=9C \rightarrow C=\frac{1}{9}$

- Para $x=1 \rightarrow 1=-2A-2B+C \rightarrow A=\frac{-1}{9}$

Por tanto: $\int \frac{dx}{x^3 - 3x^2} = \int \frac{-1}{9} \frac{1}{x} dx + \int \frac{-1}{3} \frac{1}{x^2} dx + \int \frac{1}{9} \frac{1}{x-3} dx = \frac{-1}{9} \ln|x| + \frac{1}{3x} + \frac{1}{9} \ln|x-3| + k$

m) $\int \frac{3x+\sqrt{x}}{x^2} dx = \int \left(\frac{3x}{x^2} + \frac{\sqrt{x}}{x^2} \right) dx = \int \left(\frac{3}{x} + x^{-\frac{3}{2}} \right) dx = 3 \ln|x| + \frac{x^{-\frac{1}{2}}}{-\frac{1}{2}} + k = 3 \ln|x| - \frac{2}{\sqrt{x}} + k$

n) $\int (x+3)e^{x+1} dx$. Integraremos por partes: $\begin{cases} u = x+3 & \rightarrow du = dx \\ dv = e^{x+1} dx & \rightarrow v = e^{x+1} \end{cases}$

$$\int (x+3)e^{x+1} dx = (x+3)e^{x+1} - \int e^{x+1} dx = (x+3)e^{x+1} - e^{x+1} + k = e^{x+1}(x+3-1) + k = (x+2)e^{x+1} + k$$

ñ) $\int \frac{x}{x^2 + 2x - 8} dx$. Descomponemos en fracciones simples:

$$\frac{x}{x^2 + 2x - 8} = \frac{x}{(x+4)(x-2)} = \frac{A}{x+4} + \frac{B}{x-2} = \frac{A(x-2) + B(x+4)}{(x+4)(x-2)}$$

$$\text{- Para } x=2 \rightarrow 2=6B \rightarrow B=\frac{1}{3}$$

$$\text{- Para } x=-4 \rightarrow -4=-6A \rightarrow A=\frac{2}{3}$$

$$\text{Por tanto: } \int \frac{x}{x^2 + 2x - 8} dx = \int \left(\frac{\frac{2}{3}}{x+4} + \frac{\frac{1}{3}}{x-2} \right) dx = \frac{2}{3} \ln|x+4| + \frac{1}{3} \ln|x-2| + k$$